

REGULAR ARTICLE

# Thermo-tolerance and heat shock protein of *Escherichia coli* ATCC 25922 under thermal stress using test cell method

Rui Li<sup>1</sup>, Yage Shi<sup>2</sup>, Bo Ling<sup>1</sup>, Teng Cheng<sup>1</sup>, Zhi Huang<sup>1</sup>, Shaojin Wang<sup>1,3\*</sup>

<sup>1</sup>College of Mechanical and Electronic Engineering, Northwest A&F University, Yangling 712100, China, <sup>2</sup>College of Food Science and Engineering, Northwest A&F University, Yangling 712100, China, <sup>3</sup>Department of Biological Systems Engineering, Washington State University, Pullman, WA 99164-6120, USA

## ABSTRACT

Pathogenic outbreaks related to consumption of dry foods have encouraged scientific communities to study the thermal resistance of the surrogate for thermal pasteurization validation tests. The purpose of this study was to investigate thermo-tolerance relative to mRNA levels in *E. coli* ATCC 25922 in LB broth at three temperatures after previous heat shock, and analyze heat shock protein levels after preconditioning by using aluminum test cells. The results showed that *D*-values for *E. coli* decreased as the temperature increased from 57 to 63 °C. The enhanced thermo-tolerance was found when the culture was subjected to three sub-lethal temperatures and reached maximum level at 45 °C for 5 min. The trend of relative mRNA level and the enhanced HSP 70 protein further confirmed the increased *D*-values after heat shock treatments. After storage at cold or room temperatures for a day, however, the enhanced thermo-tolerance of *E. coli* ATCC 25922 could be eliminated. Longer times at a lethal temperature are needed to guarantee the required food safety in developing effective pasteurization processes.

**Keywords:** *E. coli*; Heat shock protein; Preconditioning; Thermal processing; Thermo-tolerance

## INTRODUCTION

Outbreaks and recalls associated with pathogenic *salmonella* have been widely reported in dry foods, such as sesame seeds (Brockmann et al., 2004), spices such as paprika and pepper (Van Doren et al., 2013) and almonds (Isaacs et al., 2005). As a result, the US Department of Agriculture (USDA) regulations mandate that these agricultural products must be subjected to pasteurization processes to reduce pathogens to a reasonably acceptable level (Anon, 2007). Thermal treatments have been proposed as a promising technology for pathogen controls in postharvest agricultural products with a long and stable shelf life since they are relatively easy to apply, leave no chemical residues, and may offer some fungicidal activity (Chang et al., 2010; Liu et al., 2011).

The target pathogen cannot be directly used to conduct thermal pasteurization validation tests in food processing plants because of critical safety requirements for operators,

the product and the processing environment. Using a surrogate microorganism is an alternative way for process validations, but its thermal tolerance must be known to assure it behaves similarly to the targeted pathogen. *Escherichia coli* ATCC 25922 is nonpathogenic and has been recommended as a surrogate species of pathogenic *Salmonella* for validation of thermal pasteurization (Eblen et al., 2005). It is important to determine the thermo-tolerance of the surrogate *E. coli* for developing effective thermal treatments.

The thermo-tolerance of microorganisms is influenced by several factors, such as growth phase (Martinez et al., 2003), food composition (Juneja et al., 1998; Konate et al., 2014) and preconditioning temperatures (Juneja et al., 1998; Wiegand et al., 2009). As compared to real products, a broth culture as heating medium is widely used to determine the heat resistance of pathogens (Shenoy and Murano, 1996; Juneja et al., 2001), since it provides fast and uniform

### \*Corresponding author:

Shaojin Wang, College of Mechanical and Electronic Engineering, Northwest A&F University, Yangling 712100, China  
Tel.: +86-29 87092391; Fax: +86- 29 87091737; E-mail: shaojinwang@nwsuaf.edu.cn

**Received:** 31 July 2016; **Revised:** 10 January 2017; **Accepted:** 11 January 2017; **Published Online:** 25 January 2017

heating to achieve isothermal conditions and may represent their intrinsic heat resistance characteristics in real food (Leguerinel et al., 2007).

Thermal pasteurization protocols are developed based on the thermal resistance of pathogens under laboratory conditions. Deviation of harvest and storage conditions from room temperature prior to treatments may alter the treatment efficacy because the minimum time-temperature combinations required to reduce pathogen populations may change with pre-treatment conditions at sub-lethal warm temperatures (Juneja et al., 1998; Wiegand et al., 2009). Enhanced thermo-tolerance is related to the heat shock protein (HSP) induction (Schumann, 2003; 2007). For example, a 25 min heat shock at 46 °C for *E. coli* O157:H7 in beef gravy results in an increase in the levels of the 69 kDa DnaK and 60 kDa GroEL proteins (Juneja et al., 1998), which are classified as HSP70 and HSP60, respectively (Tobian et al., 2004). When subjected *E. coli* to 41.5 °C for 2 h heat shock, their HSP70 and HSP60 levels clearly increased (Urban-Chmiel et al., 2013). After heat shock of *Salmonella enteritidis* ATCC 4931 at 42 °C for 60 min, seven increased proteins at different molecular weights are observed (Xavier and Ingham, 1997). The relative mRNA levels and HSP have been analyzed by real-time quantitative RT-PCR (Real-time PCR), sodium dodecyl sulphate-polyacrylamide gel electrophoresis (SDS-PAGE), and Western-blot (Juneja et al., 1998; Urban-Chmiel et al., 2013). The roles of these proteins are to protect the cells against damage from high temperatures and to aid in recovery once the stress is removed (Richter et al., 2010). It is, therefore, important to determine the enhanced heat resistance of pathogens as affected by various pre-treatment conditions due to the induction and synthesis of HSP.

An aluminum test cell has been developed and applied for providing close to ideal isothermal conditions and reliable thermal death kinetic data of target pathogens (Chung et al., 2008). This test cell is designed for rapid heating of dry samples in water baths and allows easy loading and unloading of samples in a hermetically sealed cavity to evaluate the thermal resistance of vegetative cells and spores. This test cell has been successfully used to determine the *D*- and  $\alpha$ -values of *Salmonella*, *E. coli* K12, and *Clostridium sporogenes* PA 3679 spores in liquid eggs, almonds, phosphate buffer and mashed potato (Chung et al., 2008; Jin et al., 2008; Villa-Rojas et al., 2013). All of the previous research used aluminum test cells for determining the *D*- and  $\alpha$ -values of bacterial, but there is no report for determining the *D*-,  $\alpha$ -values of bacterial and relative mRNA levels of *E. coli* HSP 70 expression after previous heat shock treatment by using aluminum test cells.

The objectives of this study were to (1) determine *D*- and  $\alpha$ -values for *E. coli* ATCC 25922 in broth culture at three

temperatures using aluminum test cells, (2) obtain *D*-values and relative mRNA levels of *E. coli* HSP 70 expressions at 60 °C after heat shock at 40, 45 and 50 °C for various times, (3) analyze relative mRNA levels of heat shocked *E. coli* at 45 °C for 5 min when recovered at 4, 15 and 30 °C for various times by using real-time quantitative RT-PCR, and (4) examine the level of heat shock protein after preconditioning at 45 °C for 5 min using SDS-PAGE and Western blot analysis.

## MATERIALS AND METHODS

### Preparation of cell suspension

The material was the *Escherichia coli* ATCC 25922 strains. It was obtained from the College of Food Science and Engineering, Northwest A&F University (Yangling, China). Stock cultures were made in Luria-Bertani broth (LB: Beijing Land Bridge, Beijing, China) and stored at -20 °C with 15% (vol/vol) glycerol. Stock cultures were allowed to thaw at room temperature for 5 min. A loopful was then streaked onto LB agar and incubated at 37 °C for 24 h. A single colony was streaked onto LB agar and incubated at 37 °C for 24 h. Then, a signal colony was transferred into 30 ml LB broth and incubated for 24 h at 37 °C. Finally, 3 ml bacterial suspension was transferred to 300 ml LB broth and incubated again at 37 °C for 24 h. The cell population was adjusted to a level of 10<sup>9</sup> CFU/ml.

### Heating apparatus

To investigate the thermal resistance of *E. coli* ATCC 25922, the aluminum test cell developed at Washington State University (Chung et al., 2008) was used. This test cell consisted of a base and a screw-up cap with O-ring between the two parts for hermetical sealing. The cavity of this test cell was 18 mm in diameter and 4.5 mm height, providing 1.28 ml sample space. A pre-calibrated Type-T thermocouple (TMQSS-020-6, Omega Engineering Ltd., CT, USA) was installed in the center of the test cell to measure the core temperature of the sample, which was recorded every second by a data logger (CR1000, Campbell Scientific Inc., Logan, Utah, USA). The time/temperature history was measured for each run using a non-inoculated sample (Luria-Bertani broth), and recorded in a computer. The recorded temperature-time history was used to determine the come-up time for each of three target temperatures. Before each use, the test cells were sanitized according to Jin et al., (2008).

### Effects of experimental conditions on the microbial thermo-tolerance

#### Heat treatment

One ml of *E. coli* ATCC 25922 bacterial suspensions with 10<sup>9</sup> CFU/ml cell populations was placed inside each test cell. Hermetically sealed test cells were then submerged completely in a water bath (SC-15, Scientz Biotechnology,

Ningbo, China). The temperature of the water bath was controlled at 57, 60 or 63 °C. The come-up time for the test cell core to reach within 0.5 °C of each set-point temperature was determined and used as time zero to provide close-to-ideal isothermal conditions. Test cells were removed at five different time intervals, depending on the temperature, to achieve at least 5-log reductions. After holding, the test cells were immediately placed in an ice-water bath until further analysis was performed. As controls, samples in the test cells were treated in the same way for the longest holding time without heating. All the experiments were repeated three times. The semi-log plot of microbial survival curves was used to determine *D*-value (decimal reduction time) and  $\alpha$ -value (temperature changes needed for 90% reduction in *D*-values).

### Heat shock

One ml of *E. coli* ATCC 25922 bacterial suspensions was transferred to each test cell, then heat shocked at 40, 45 or 50 °C temperature controlled water bath for 2, 5, 8 or 11 min, and the total test combinations were 12 treatment conditions. After heat shocking, test cells were immediately cooled in ice water. Thereafter, test cells were then submerged completely in a water bath at 60 °C. After holding for 0-5 min, duplicate test cells were removed from the water bath and immediately immersed in an ice-water bath until further analysis was performed. A sample from the test cell that was not exposed to heat shock served as an untreated control.

### Recovery tests

Test cells containing 1 ml of *E. coli* ATCC 25922 bacterial suspensions were sealed and then heat shocked at 45 °C for 5 min, and subsequently stored at 4, 15 or 30 °C for 4, 8, 14, 24 or 32 h, all of these stored conditions were 15 combinations test. After each storage period, real-time PCR was conducted to detect the relative expression levels of *E. coli* ATCC 25922 mRNA.

### Microbiological enumeration

One ml of *E. coli* ATCC 25922 bacterial suspensions was taken from test cells and then washed by using 1 ml sterile physiologic saline. Subsequent 10-fold serial dilutions were performed in 4.5 ml of sterile physiologic saline; 100 µl of each one was spread onto duplicated LB agar and incubated at 37 °C for 24 h, and cultures were counted to enumerate the plates with 30-300 colonies.

### RNA isolation, reverse transcription and real-time PCR

After heat shock at temperatures of 40, 45 and 50 °C for various times, the total RNA from the cell suspension was extracted by using RNAiso Plus (TaKaRa, Dalian, China) following the manufacturer's instructions. The RNA concentration and purity were determined from measuring the optical density (OD) at 260 and 280 nm wavelengths using an Epoch microplate spectrophotometer

(BioTek Instruments. Inc., USA). The OD<sub>260/280</sub> ratios were from 1.8 to 2.1 for all samples. A 500 ng total RNA template was used to convert mRNA into cDNA using a Prime Script™ RT Master Mix (TaKaRa, Dalian, China) according to the manufacturer's instructions. Real-time PCR was then performed using a CFX Connect™ Real-Time PCR Detection System (Bio-Rad, CA, USA). For normalization, 16s-rRNA was employed. Each experiment was repeated independently at least three times, and the fold change in the expression of DnaK gene was analyzed via a  $2^{-\Delta\Delta C_t}$  method (Schmittgen and Livak, 2008). The primers for real-time PCR corresponding to each gene were designed using Primer 5.0 software and listed in Table 1.

In a series of recovery experiments, *E. coli* ATCC 25922 was heated at 45 °C for 5 min and subsequently stored at 4, 15 or 30 °C for 4, 8, 14, 24 or 32 h. After each storage period, Real-time PCR was conducted to detect the relative expression levels of *E. coli* ATCC 25922 mRNA. For protein analysis, 1 ml aliquots from each sample were collected and centrifuged; 1 ml aliquots from cells that had not been heat-shocked were also harvested to serve as a negative control. The cells were re-suspended in 200 µl of SDS-PAGE sample buffer, boiled for 5 min, and then carried out in a 5–10% polyacrylamide gel gradient in standard Tris/glycine chamber buffer (0.025 M Tris/HCl, 0.129 M glycine, 0.1% SDS) at 100 V in a Mini-Protean II apparatus (Bio-Rad). Broad range (6.5–200 kDa; Bio-Rad) was used as a molecular mass standard. Pictures were taken using a digital camera and the bands determined by comparison with molecular weight of the standards.

### Western blot analysis

Western blotting with mouse anti-Hsp70 (LSBio) mAbs was used for identified protein fractions in cell lysates. The protein fractions obtained in electrophoresis were transferred to a 0.2 mm nitrocellulose membrane (Bio-Rad), described by Towbin et al., (1979). The membranes were incubated in blocking buffer with 5% non-fat milk for 1 h at room temperature, then washed twice in TBST buffer (TBS with 0.05% Tween 20, pH 7.5). The membranes were then incubated at 4 °C in mouse anti-Hsp70 overnight, washed with TBST and incubated for 60 min at room temperature with horseradish-peroxidase-conjugated mouse anti-IgG

**Table 1: Sequences of primers for real-time PCR used in this study**

| Gene    | Sense<br>antisense | Primer sequence       | Size (bp) |
|---------|--------------------|-----------------------|-----------|
| DnaK    | Sense              | TCTGGTTGGTCAGCCGGCTAA | 158       |
|         | Antisense          | CGTCGCCGTTATCAGCAGCAA |           |
| 16srRNA | Sense              | TGCCATCGGATGTGCCAGAT  | 173       |
|         | Antisense          | CATCAGGCTTGCGCCATTGT  |           |

All primers were synthesized by Invitrogen Biotech Co., Ltd. (Beijing, China)

antibodies (Biokom). The membranes were stained using a color development kit (Bio-Rad). Analysis was conducted using Quantity One software (Bio-Rad).

**Statistical analysis**

*D*-values and relative mRNA level of DnaK expressions in *E. coli* ATCC 25922 from heat shock cells subjected to heat treatment with three independently replicated experiments were determined. Data were analyzed using the Statistical Product and Service Solutions (SPSS statistical software version 17.0). The Duncan multiple range test was used as post hoc determination of significant differences ( $P < 0.05$ ). The *Z*-values were determined from the reciprocal of the slope obtained by plotting the log of the *D*-values against exposure temperatures.

**RESULTS AND DISCUSSION**


**Heat resistance of *E. coli* at three temperatures**

A typical temperature-time history and survival curves for *E. coli* ATCC 25922 in test cell subjected to a water bath at 57, 60, and 63 °C are shown in Fig. 1. The sample core temperatures reached the set-point quickly, resulting in the similar come-up times of 20 s for three temperatures and providing close to ideal isothermal conditions. This come-up time was similar to that of mashed potato in a capillary tube (Chung et al., 2008), but shorter than those of mashed potato in glass tubes and

of almond kernels in the same cell (Villa-Rojas et al., 2013) caused by poor heat conduction in solid materials. After short coming up times, the initial reduction of *E. coli* was negligible based on small intercepts of the linear regression equations. The time to reduce 4-log populations was 4.5, 2.8 and 1.5 min at 57, 60 and 63 °C, respectively, indicating that *D*-values of *E. coli* ATCC 25922 decreased as the temperature increased from 57 to 63 °C, resulting in *z*-value of 13.3 °C. This decreasing trend of *D*-values has been observed in other studies (Rajkowski, 2012).

**Effect of heat shock on the heat resistance of *E. coli***

The *D*-values and relative mRNA level of DnaK expression of *E. coli* ATCC 25922 at 60 °C after heat shocking at three given temperature/time combinations are listed in Table 2. The results showed that the maximum *D*-value appeared after heat shocking at 45 °C for 5 min. The significant difference in *D*-value ( $p < 0.05$ ) was observed between 40 and 45°C for heating 5 min, between 45 and 50°C for heating 2 and 8 min, or between 5 min heating and other durations at 45°C. The enhanced thermo-tolerance was also found in *E. coli* O157:H7 in trypticase soy broth, *L. monocytogenes* Scott A in trypticase soy + 0.6 % yeast extract broth and *E. coli* ATCC 25922 in nutrient broth after heat shock treatments (Gadzella and Ingham, 1994; Linton et al., 1992; Murano and Pierson, 1992; Yuk and Marshall, 2003). From a food safety standpoint, therefore,



**Fig 1.** A typical temperature-time history (a) and survival curves (b) for *E. coli* ATCC 25922 in the test cell center subjected to a water bath at  $\Delta$ 57,  $\square$ 60, and  $\circ$ 63 °C.

**Table 2:** *D*-values (Mean±SD) and relative mRNA level of DnaK expressions (Mean±SD) of *Escherichia coli* ATCC 25922 at 60°C after heat shock at three given temperatures for various times

| Duration of heat shock (min) | <i>D</i> -value (min) after following heat shock temperatures ( °C) |             |             | mRNA levels after following heat shock temperatures ( °C) |              |              |
|------------------------------|---|-------------|-------------|---|--------------|--------------|
|                              | 40  | 45          | 50          | 40  | 45           | 50           |
| 0 <sup>#</sup>               | 0.78±0.01aA*  | 0.77±0.01aA | 0.79±0.01aA | 0.489±0.03aA*   | 0.492±0.03aA | 0.490±0.03aA |
| 2                            | 0.91±0.02bA   | 0.92±0.02bA | 1.03±0.02cB | 0.581±0.03bA  | 0.798±0.02bB | 4.805±0.06bC |
| 5                            | 0.92±0.02bA   | 1.25±0.03cB | 1.19±0.03dB | 0.795±0.02cA  | 16.91±0.05cB | 8.406±0.13cC |
| 8                            | 1.01±0.03cAB  | 0.96±0.02bA | 1.05±0.02cB | 1.226±0.04dA  | 2.947±0.05dB | 5.048±0.08dC |
| 11                           | 0.95±0.04bcA  | 0.94±0.02bA | 0.91±0.04bA | 1.048±0.02eA  | 1.976±0.03eB | 0.593±0.03eC |

<sup>#</sup>0 means non-heat shock treatment and was used as a control group. \*Different lower and upper case letters indicate that means are significantly different at  $P=0.05$  among duration of heat shock and heat shock temperatures, respectively



heat-shocked cells need to be heated for longer time or higher temperature to achieve the same lethal level required by non-heat-shocked cells (Farber and Pagotto, 1992). The trend of relative mRNA level as a function of heat shock temperature and time was observed to be similar to that of *D*-values. The highest relative mRNA level of HSP 70 expressions of *E. coli* ATCC 25922 was also observed at 45 °C for 5 min. This indicated that after heat stress, the relative mRNA level of HSP 70 expression increased, since high temperatures might enhance the activity of transcription factors, thus strengthen the synthesis of the relative mRNA level of HSP 70 expression (Craig et al., 1991; Gabai et al., 1997).


### Recovery characteristics of *E. coli* after removing heat shock

Relative mRNA levels of DnaK expression of *E. coli* ATCC 25922 when stored at 4, 15 and 30 °C for various times after heat shocked at 45 °C for 5 min shown in Fig. 2, which was the most heat resistant conditions. The results show that the relative mRNA level decreased after 4 h storage at 30 °C but 8 h at 4 °C and 15 °C and reached the level below the control after 24 h. After storage at cold or room temperatures for a day, the enhanced thermo-tolerance of *E. coli* ATCC 25922 could be eliminated since the relative mRNA levels dropped down to less than that of controls. This study underlines the importance of keeping contaminated agricultural products in a room temperature condition for a period of time to avoid pretreatment conditioning. These findings are in agreement with that reported by Jørgensen et al., (1999) who indicated that heat-shock-induced thermo-tolerance of *L. monocytogenes* in tryptic phosphate broth was maintained for a longer time at 4 and 10 °C than stored at 30 °C. However, the shorter storage duration at higher temperatures to reduce the heat shock proteins is different from that found by Juneja et al., (1998) and Auffray et al., (1995) in which *E. coli* O157:H7 cells in beef lost their thermo-tolerance after 14 h at 4 °C and after 24 h in beef at 15 or 28 °C. The persistence of heat-shock-induced thermo-tolerance may be due to different metabolic states of the tested cells (Lindquist, 1986).


### Levels of heat shock protein synthesis

To determine whether the heat shock treatment used in the present study resulted in an increase in the levels of any major heat-shock protein (Hsp), the protein patterns of *E. coli* ATCC 25922 both in control cells and heat shocking at 45 °C for 5 min were compared following electrophoresis. Because heat-inducible members of HSP 70 family (DnaK) play an important role in thermo-tolerance (Schumann, 2003) meanwhile DnaK can prevent protein aggregation and promote renaturation at high temperatures (Isaacs et al., 2005). Thus in our

experiment thermo-tolerance of *E. coli* ATCC 25922 after heat shock accompanied by the induction of HSP 70 was determined. (Fig. 3a) shows that the level of one protein with an apparent molecular mass of approximately 69 kDa was found both in heat-shocked and control cells, but the level of 69 kDa protein in heat-shocked cells was



**Fig 2.** Relative mRNA levels of DnaK expressions of *E. coli* ATCC 25922 when stored at 4, 15 and 30 °C for various times after heat shocked at 45 °C for 5 min. Error bars show the standard error of the means. Real-time PCR data were obtained as Ct values and amounts of DnaK mRNA were normalized to 16s-rRNA. Control means after heat shocked at 45 °C for 5 min. Different letters indicate that means in control are significantly different at  $P=0.05$  from treatments.



**Fig 3.** SDS-PAGE protein profiles of the *E. coli* ATCC 25922 strains (a) and Western-blot reactions of *E. coli* ATCC 25922 with mouse monoclonal anti-HSP70 antibodies (b). Lane 1 represents protein samples from the control, non-heat shocked cells; lane 2 represents protein samples from cells heat-shocked at 45 °C. 10 ul proteins loaded in each well to determine the HSP protein.

slightly higher than that in non-heat-shocked cells. To determine whether this protein detected by SDS-PAGE analysis corresponded to the HSP 70, a Western-blot was performed using monoclonal antibodies specific for *E. coli* HSP 70 (69 kDa). This protein was detected on the blot in control cells, but after heat shocking at 45 °C for 5 min treatment, the level of this protein increased, determined by densitometry (Fig. 3b).

The observed HSP 70 protein in controls was probably caused by many diverse heat shock proteins reported in *E. coli* (Neidhardt et al., 1984). However, the increased level of HSP 70 protein after heat shock was similar to results obtained by Auffray et al., (1995) and Juneja et al., (1998). This is because the loss of normal cellular functions due to sudden temperature increase is recovered by the synthesis of stress proteins such as Hsps (Juneja et al., 1998; Li and Mak, 1989; Urban-Chmiel et al., 2013).

High temperatures are stress conditions that induce responses in an organism to cope with these stresses. One of the most studied responses is the induction of heat shock proteins in pathogens. The major inducible HSP70 may reduce the inactivation level required by food safety. Determining the heat-shock response and thermo-tolerance is practically important to control *E. coli* ATCC 25922 in agricultural products that are heated under different harvesting, processing and storage conditions. Therefore, the enhanced heat resistance of *E. coli* ATCC 25922 due to heat shock must be considered while designing effective thermal processes.

## CONCLUSIONS

In this study, the heat resistance of *E. coli* ATCC 25922 in LB broth at three temperatures was obtained with test cell methods. The preheating conditions at three sub-lethal temperatures resulted in increased heat resistances of *E. coli* with maximum *D*-values after heat shock at 45 °C for 5 min. The trend of relative mRNA level after heat shock treatment was similar to that of *D*-values. The increased heat resistance could be eliminated by adding one-day storage of the contaminated products at room or cold temperatures. Avoiding heat shock proteins or increasing temperatures or treatment times must be considered to ensure food safety while designing effective thermal processes to control *E. coli* ATCC 25922 in postharvest agricultural products. Further experiments should be conducted to study the thermo-tolerance of *E. coli* ATCC 25922 in real foods and validate the results obtained in this study.

## ACKNOWLEDGEMENTS

This research was conducted in the College of Mechanical and Electronic Engineering, Northwest A&F University, and supported by research grants from General Program of National Natural Science Foundation of China (31371853). The authors thank Kexin Gao for her help on RT-PCR evaluation and Lijun An, Xiayan Liu and Yuanfeng Li for their helps on the SDS-PAGE and Western-blot analyses.

## Author contributions

Conception design: S.W. Performing experiment and data analysis, drafting of the manuscript: R.L. Assist Experiment and critical revision of the manuscript: Y.S, B.L, T. C, and Z. H. All authors reviewed the manuscript.

## REFERENCES

- Anon. 2007. Almonds Grown in California; Outgoing Quality Control Requirements. Federal Register 72, 15021-15036. Agricultural Marketing Service, USDA.
- Auffray, Y., E. Lecesne, A. Hartke and P. Boutibonnes. 1995. Basic features of the *Streptococcus thermophilus* heat shock response. *Curr. Microbiol.* 30: 87-91.
- Brockmann, S. O., I. Piechotowski and P. Kimmig. 2004. *Salmonella* in sesame seed products. *J. Food Protect.* 67: 178-180.
- Chang, S. S., A. R. Han, J. I. Reyes-De-Corcuera, J. R. Powers and D. H. Kang. 2010. Evaluation of steam pasteurization in controlling salmonella serotype enteritidis on raw almond surfaces. *Lett. Appl. Microbiol.* 50: 393-398.
- Chung, H. J., S. L. Birla and J. Tang. 2008. Performance evaluation of aluminum test cell designed for determining the heat resistance of bacterial spores in foods. *LWT - Food Sci. Technol.* 41: 1351-1359.
- Craig, E. and C. A. Cross. 1991. Is hsp70 the cellular thermometer. *Trends Biochem. Sci.* 16: 35-40.
- Eblen, D. R., B. A. Annous and G. M. Sapers. 2005. Studies to select appropriate nonpathogenic surrogate *Escherichia coli* strains for potential use in place of *Escherichia coli* O157: H7 and *Salmonella* in pilot plant studies. *J. Food Protect.* 68: 282-291.
- Farber, J. M. and F. Pagotto. 1992. The effect of acid shock on the heat resistance of *Listeria monocytogenes*. *Lett. Appl. Microbiol.* 15: 197-201.
- Gabai, V. L., A. B. Meriin, D. D. Mosser, A. W. Caron, S. Rits, V. I. Shifrin and M. Y. Sherman. 1997. HSP70 prevents activation of stress kinase – A novel pathway of cellular thermotolerance. *J. Biol. Chem.* 272: 18033-18037.
- Gadzella, T. A. and S. C. Ingham. 1994. Heat-shock, anaerobic jar incubation and fluid thioglycolate medium have contrasting effects on D-Values of *Escherichia-coli*. *J. Food Protect.* 57: 671-673.
- Isaacs, S., J. Aramini, B. Ciebin, J. A. Farrar, R. Ahmed, D. Middleton, A. U. Chandran, L. J. Harris, M. Howes, E. Chan, A. S. Pichette, K. Campbell, A. Gupta, L. Y. Lior, M. Pearce, C. Clark, F. Rodgers, F. Jamieson, I. Brophy and A. Ellis. 2005. An international outbreak of salmonellosis associated with raw almonds contaminated with a rare phage type of *Salmonella*

- enteritidis*. J. Food Protect. 68: 191-198.
- Jin, T., H. Zhang, G. Boyd and J. Tang. 2008. Thermal resistance of *Salmonella enteritidis* and *Escherichia coli* K12 in liquid egg determined by thermal-death-time disks. J. Food Eng. 84: 608-614.
- Jørgensen, F., T. B. Hansen and S. Knøchel. 1999. Heat shock-induced thermotolerance in *Listeria monocytogenes* 13-249 is dependent on growth phase, pH and lactic acid. Food Microbiol. 16: 185-194.
- Juneja, V. K., B. S. Eblen and G. M. Ransom. 2001. Thermal inactivation of *Salmonella spp.* In chicken broth, beef, pork, turkey, and chicken: Determination of D- and Z-values. J. Food Sci. 66: 146-152.
- Juneja, V. K., P. G. Klein and B. S. Marmer. 1998. Heat shock and thermotolerance of *Escherichia coli* O157:H7 in a model beef gravy system and ground beef. J. Appl Microbiol. 84: 677-684.
- Konate, M., E. E. Akpa, L. B. Koffi, K. A. S. Kra, R. M. Megnanou and S. Niamke. 2014. Isolation of thermotolerant and high acetic acid-producing *Acetobacter pasteurianus* from Ivorian palm wine. Emirates J. Food Agric. 26: 773-785.
- Leguerinel, I., I. Spegagne, O. Couvert, L. Coroller and P. Mafart. 2007. Quantifying the effects of heating temperature, and combined effects of heating medium pH and recovery medium pH on the heat resistance of *Salmonella typhimurium*. Int. J. Food Microbiol. 116: 88-95.
- Li, G. C. and J. K. Mak. 1989. Re-induction of hsp70 synthesis: An assay for thermotolerance. Int. J. Hyperthermia. 5: 389-403.
- Lindquist, S. 1986. The heat-shock response. Annu. Rev. Biochem. 55: 1151-1191.
- Linton, R. H., J. B. Webster, M. D. Pierson, J. R. Bishop and C. R. Hackney. 1992. The effect of sublethal heat-shock and growth atmosphere on the heat-resistance of *Listeria-Monocytogenes Scott-A*. J. Food Protect. 55: 84-87.
- Liu, Y. H., J. Tang, Z. H. Mao, J. H. Mah, S. S. Jiao and S. J. Wang. 2011. Quality and mold control of enriched white bread by combined radio frequency and hot air treatment. J. Food Eng. 104: 492-498.
- Martinez, S., M. Lopez and A. Bernardo. 2003. Thermal inactivation of *Enterococcus faecium*: Effect of growth temperature and physiological state of microbial cells. Lett. Appl. Microbiol. 37: 475-481.
- Murano, E. A. and M. D. Pierson. 1992. Effect of heat-shock and growth atmosphere on the heat-resistance of *Escherichia-coli* O157-H7. J. Food Protect. 55: 171-175.
- Neidhardt, F. C., R. A. VanBogelen and V. Vaughn. 1984. The genetics and regulation of heat-shock proteins. Annu. Rev. Genet. 18: 295-329.
- Rajkowski, K. T. 2012. Thermal inactivation of *Escherichia coli* O157:H7 and *Salmonella* on catfish and tilapia. Food Microbiol. 30: 427-431.
- Richter, K., M. Haslbeck and J. Buchner. 2010. The heat shock response: Life on the verge of death. Mol. Cell. 40: 253-266.
- Schmittgen, T. D. and K. J. Livak. 2008. Analyzing real-time PCR data by the comparative C-T method. Nat. Protoc. 3: 1101-1108.
- Schumann, W. 2003. The *bacillus subtilis* heat shock stimulon. Cell Stress Chaperones. 8: 207-217.
- Schumann, W. 2007. Thermosensors in eubacteria: Role and evolution. J. Biosci. 32: 549-557.
- Shenoy, K. and E. A. Murano. 1996. Effect of heat shock on the thermotolerance and protein composition of *Yersinia enterocolitica* in brain heart infusion broth and ground pork. J. Food Protect. 59: 360-364.
- Tobian, A. A. R., D. H. Canaday and C. V. Harding. 2004. Bacterial heat shock proteins enhance class II MHC antigen processing and presentation of chaperoned peptides to CD4(+) T cells. J. Immunol. 173: 5130-5137.
- Towbin, H., T. Staehelin and J. Gordon. 1979. Electrophoretic transfer of proteins from polyacrylamide gels to nitrocellulose sheets: Procedure and some applications. Proc. Natl. Acad. Sci. USA. 76: 4350-4354.
- Urban-Chmiel, R., M. Dec, A. Puchalski and A. Wernicki. 2013. Characterization of heat-shock proteins in *Escherichia coli* strains under thermal stress *in vitro*. J. Med. Microbiol. 62: 1897-1901.
- Van Doren, J. M., K. P. Neil, M. Parish, L. Gieraltowski, L. H. Gould and K. L. Gombas. 2013. Foodborne illness outbreaks from microbial contaminants in spices, 1973-2010. Food Microbiol. 36: 456-464.
- Villa-Rojas, R., J. Tang, S. Wang, M. Gao, D. H. Kang, J. H. Mah, P. Gray, M. E. Sosa-Morales and A. Lopez-Malo. 2013. Thermal inactivation of *Salmonella enteritidis* PT 30 in almond kernels as influenced by water activity. J. Food Protect. 76: 26-32.
- Wiegand, K. M., S. C. Ingham and B. H. Ingham. 2009. Survival of *Escherichia coli* O157:H7 in ground beef after sublethal heat shock and subsequent isothermal cooking. J. Food Protect. 72: 1727-1731.
- Xavier, I. J. and S. C. Ingham. 1997. Increased D-values for *Salmonella enteritidis* following heat shock. J. Food Protect. 60: 181-184.
- Yuk, H. G. and D. L. Marshall. 2003. Heat adaptation alters *Escherichia coli* O157:H7 membrane lipid composition and verotoxin production. Appl. Environ. Microbiol. 69: 5115-5119.